

Effect of Infection by Entamoeba Histolytica on Some Blood Parameters in Kufa City

Kareem A. Hammad¹
kareem.hammady@hiuc.edu.iq

Maysoon AL-Adraawy²
kin.msn@atu.edu.iq

Haqi A .A .Al-Fatlway³
kin.hke@atu.edu.iq

Alaa Kareem Mohammed⁴
alla.kareem@hiuc.edu.iq

Esam M.Turky⁵
kin.asm@atu.edu.iq

Abstract: This study was carried out to study the effect of Entamoeba histolytica infection and its impact on some blood parameter such as hemoglobin, packed cell volume (p.c.v.), cholesterol and Triglyceride (TG) among 430 patients who attended AL-Furat AL-owset Teaching hospital in Kufa city / Iraq during six month period from March and September 2018. Those patients suffered from diarrhea and abdominal discomfort. The frequency of Entamoeba histolytica occurring was (70) infection out of (430, 16.3%). Among those (70) infected there were male n=32/70,45.7%, while females represented n=38/70,54.3%. The result of this study showed no significant affect in cholesterol and Triglyceride but significant decrease in Hb and pcv in comparison with control group of (30) healthy individual which was included in this study.

Keywords: Entamoeba histolytica, cholesterol, Iraq, Blood parameter.

¹ Assist. Prof. Dr.: Medical Laboratory Techniques Department, Al-Hikma University College, Baghdad, Iraq

² Assist. Prof.: Kufa Technical Institute, ALFurat AL Awsat Technical University, Iraq

³ Assist. Prof.: Kufa Technical Institute, ALFurat AL Awsat Technical University, Iraq

⁴ Prof. Dr.: Medical Laboratory Techniques Department, Al-Hikma University College, Baghdad, Iraq

⁵ Assist. Prof.: Kufa Technical Institute, ALFurat AL Awsat Technical University, Iraq

1. Introduction

Entamoeba histolytica among the most common protozoan parasite which causes Amoebiasis [1]. E. histolytica is the most common intestinal parasite of human which affects the large intestinal area causing ulceration and spread dysentery worldwide [2]. It is a unicellular protozoan parasite with mortality rate over 100000 worldwide yearly [3]. About 10% of cases are asymptomatic, although, the parasite can invade the gut wall causing severe ulceration and amoebic dysentery characterized by blood [4].

The parasite transmission through contaminated food and water via fecal-oral route [5]. This parasite causing intestinal colic, diarrhea occasionally diarrhea with bleeding of mucosa and loosen of appetite, vomiting, loss of Na⁺, K⁺ which may lead to malnutrition and anemia [6]. This parasite infection could be transmitted to extra intestinal region such as liver and causes abscesses [7].

The aim of this study is to detect the parasite occurrence in this city within certain time of six months and to measure the impact of infection on blood parameter such as Hb, pcv, cholesterol, Triglyceride in compared with control group in order to pay attention on the Health problems as well as the disease, we associated with this parasite.

2. Materials and Methods

2.1 Defection of the parasite

Stool samples were collected from a total of (430) patients included males and females with different ages (1 years to 65 years old). These patients attended AL-Furat AL Owset teaching hospital within six months period from March September 2018. The stool samples were collected in clean plastic bottles labeled with specific patients data i.e. name, age, sex. These samples were examined under microscope after mixing with normal saline during slide preparation. Out of (430) stool samples only (70) stool samples confirmed cyst E. histolytica. Those samples were diagnosed on the base of clinical finding and microscopically stool examination according to standard laboratory procedure.

2.2 Measurement of blood parameter

The blood parameters included the measurement of Hb %, by using cyan methemoglobin method according to Salh's method, in this method the Drabkin solution was used for dilution which contains cyanide ferric potassium $K_3Fe(CN)_6$ and potassium cyanide (KCN) these two compound associated forming complex compound of melhemoglobin. The color intensity was measured at (540) wavelength by using spectrophotometer. As for pcv measured by using

capillary test tubes, micro centrifuge and hematocrit reader to measure and determine the volume of packed RBCs cells [8].

2.3 Estimation of total serum cholesterol

This was done by enzymatic and colorimetric method in which cholesterol esterolysis to cholesterol and fatty acid by cholesterol esterase [9].

3. Results and Discussion

The infection with *E. histolytica* was significantly high the frequency of infection was (16.3%). The male represented (45.7%) percentage while it was for female (54.3%). These results showed in Table 1 the distribution according to the gender.

Table 1: The distribution of *E. histolytica* infections

Sex	Number	Percentage
Male	32	45.7%
Female	38	54.3%
Total	70	100%

There was significant difference according to gender $p > 0.05$. These results were in agreement with previous study was done by Wassir (2010) [11]. The distribution of infection with *E. histolytica* according to the months of period of this study the highest number was in July, August both months represented the showed the number test in Iraq. The results in Table 2 decline in number of infections started May, April and March as the temperature started dropped in these three months of this study.

Table 2: The distribution of *E. histolytica* infections according to months of study

Months	No. of samples	Number of positive infections	Percentage %	Male	Female
August	94	16	17	5	11
July	102	18	17.6	11	7
June	98	15	15.3	7	8
May	58	13	46.5	4	9
April	43	6	2.3	3	3
March	35	2	1.3	2	0
Total	430	70	100	32	38

The result showed in table (2) was in agreement with previous study was done by [12].The result for serum cholesterol estimation as shown in figure (1) showed significant decrease $p < 0.05$ in cholesterol concentration among patients which was $147.2 \mu\text{g/dl}$ in comparison with control group which as $151.9 \mu\text{g/d}$. Figure (1)




Figure 1: Cholesterol concentration among patients and control group

There were significant differences among control group and patients ($P < 0.05$) The results in Figure 2 showed there were significant decrease ($p < 0.05$) for serum Triglyceride concentration which in patients 116.1 mg/dl while its concentration for control groups 120.6 mg/dl .




Figure 2: Serum Triglyceride concentration among patients and control group

The results in Figure 1 and Figure 2 showed significant decline in cholesterol and Triglyceride respectively among infected patients with *E. histolytica* and control group. This may be due to the virulence and pathogenicity of adhesive on the epithelial cells [11]. This parasite which causes amoebiasis which effect by reducing the cholesterol absorption in related to liver abscess this causes the cholesterol and triglyceride to be depicted from serum [12].

The previous studies [13, 14] showed that this parasite become more virulent in the presence of cholesterol .Also *E. histolytica* utilized cholesterol for their growth [15], as well as the using of cholesterol with host factor such as normal flora may be needed in encystations of *E. histolytica* [16].

The measurement of Hb% and PCV showed reduction only in (No. 56) which represented (56/430, 13%). The results of this study showed increased of incidence of *E. histolytica* infection during August ,July, June in comparison with May , April and March as the Temperature decline this may be in related for encystation of *E. histolytica* as well as decrease food hygiene during summer months . These results were in agreement with pervious study [17, 18, 19].

The results of this study showed increased of infection among females more than in males this may be related to physiological behavioral as well as other difference [20]. The results in related to Hb% and Triglyceride deficiency may be to loss of blood which associated with bleeding through amoebiasis dysentery and ulceration [21].

4. Conclusion

This study showed that there is a need for more study and investigation of *E. histolytica* occurrence for other places of the country and apply notional medical health strategy for limitation for *E. histolytica* infection among population.

References

- [1] Aque R, Hnston C.D, Hughes M , Houpt E , Petriew A. ,(2003)."Amebiosis" N Engl J media , 348: 1565 – 73
- [2] Ryan, K. J. and Ray, C.g. (2004) . sherries medical microbiology, 4th Edition, Mc graw Hill, New York, 733-778.
- [3] Fotedar, R. , stark , Beebe , N., Marriott, D., Etlis, J. and Horkness , J. (2007))"laboratory diagnostic techniques for Entamoeba species " clinical microbiology Reviews 20: 511-532,
- [4] G.ML and Taunich , Elongitudinal (2003)." study of intestinal Entamoeba histolytica infection in a symptomatic adult carrier s .Journal of clinical microbiology 41: 4745.4750,

- [5] Lucas, R. and Uporoft, J.A.(2001) " Clinical significance of the redefinition of the agent of amoebiasis " *Revista latino. American de microbiology* 43: 183-187.
- [6] Keen, E.C.(2013)*Beyon phage Therapy : virotherpy of protozoal Diseases . Future microbiology* , 81821-823.
- [7] WHO(1997).WHO, PAHO, Un, EscoReprti a A constitution with Experts on Amoebiasis Mexico city ,Mexico , 28-29 January 1997. *Epidemiological Bulletin*, 18,13-14.
- [8] Sood. R.(1994) *practical pathology and microbiology JAPEE Brether*, New Delhis , 165-188.
- [9] Allain, A.C.; Poor ,L.S. and chan , c.s.g. (1997)."Enzymatic determination of total serum cholesterol" *Clin.chom.*:20-47.
- [10] Fossati, P. and prencipe, L.(1982) " serum Tg, determination colormeterically with one enzyme that produce inflammatory reaction " *Amj. Patholo*; 107-397.
- [11] Gareia , A.F.;Chang , T.H.,Benchimol , Mi; Klump A,D.J.;lehker., M.W. and Aldeverle, J.F.(2003)." Iran and contact with host cell induce expression of adhesion on surface of T.vaginali,Nol. *Microbial.*,47: 1207-1224.
- [12] D.bonsal,H.S.bhatti and R.sehgal,(2005)."Altered lipid porameters in patient effected with Entamoeba histolytica. Entameba disper and Giardia lamblia" *British Journal of biomediral , science ,india* , , 62(2).
- [13] Sharma R., 1959." Effect of cholesterol on the growth and virulence of Entamoeba histolytica". *Trans Royoctrop and Med. Hyy*;53:278.
- [14] Sigh BN, srivastora RVN oputta gp.(1971) " virulence of strain of Entamoeba histolytica torats and the effect of cholesterol rutcaecal and hamster liver passageon the virulence of non in vassivestrions " *Indian J.Exp. Biol .* , 4:21.
- [15] Kaul D, Rani R., Sehgal R .(2001). " Receptor, Cle- regulates Giardia encystation processes (2001). " *molcell Rioch emi*,225: 167- 9.
- [16] Byers J.Eaigle W, Eichinger D. (2005)"colonic short chain fatt acids inhibit enystation of Entamoeba invoders" . *cell microbial*; 7: 269 – 279.
- [17] Bazzaz, A.A. and Ahmed , N.A. (2006). Prevalence of some parasite infection disease within tarkuk city for year 2009 – 2014. *Europeon Journal of pharmaceutical and medical Resorch* , 3: 13-19.
- [18] Nassir,A.M.A.(2010)." Diagnosis Epidemiological study of some species of Entamoeba using PCR in Tikrit.MCS Thesis , foruly of science , university of Tikrit .

- [19] AL Musawi, A,A,H, AL Yozi , W.jm, and ALmauooli H. K. (2006). Studies on porusite Diarrhea in children at Karbala city, scientific Journal of Karbala university , 4: 17-30.
- [20] Zahida , T. Shebona , K and Lashari , M. H. (2010). " Prevelance of Entamoeba histolytica in Human , Pakistani Journal of phormcentical science , 23: 344- 348.
- [21] Farror , J.Halez , p. ,Junghons, T. Kang, glalloo, Dand white , N. J. (2013). " manson, Tropical Diseases Elsevier. Health sciences , 664-671.

تأثير الاصابة بالاميبا الحالة للنسيج على بعض معايير الدم في مدينة الكوفة

حقي عبدالعباس عيسى³

kin.hke@atu.edu.iq

ميسون خضير عبدالعباس²

kin.msn@atu.edu.iq

كريم عليوي حمادي¹

kareem.hammady@hiuc.edu.iq

عصام محمد تركي⁵

kin.asm@atu.edu.iq

علاء كريم محمد⁴

alla.kareem@hiuc.edu.iq

المستخلص: أجريت هذه الدراسة لمعرفة تأثير الاصابة بطفيلي *Entamoeba histolytica* وتأثير هذه الاصابة على بعض معايير الدم مثل الهيموكلوبين ، p.c.v الكولستيرول والدهون الثلاثية بين 430 مريض وفدوا الى مستشفى الفرات الاوسط التعليمي في مدينة الكوفة في العراق . لفترة ستة أشهر من شهر آذار ولغاية شهر ايلول 2018 . هؤلاء المرضى كانوا يعانون من حالات اسهال والالام البطنية. كانت نسبة الاصابة 16.3% حيث بلغ عدد المصابين (70) مصاب حيث اصابة الرجال 45.7%(70/32) ونسبة الاناث 54.3% (70/38) أظهرت الدراسة نقص معنوي في مستوى الهيموكلوبين و pcv مقارنة بمجموعة السيطرة البالغة 30 فرد من الاصحاء .

الكلمات المفتاحية: الاميبا الحالة للنسيج، الكولستيرول، العراق، معايير الدم

¹ استاذ مساعد دكتور: قسم تقنيات المختبرات الطبية – كلية الحكمة الجامعة – بغداد - العراق

² استاذ مساعد: المعهد التقني – كوفة – جامعة الفرات الاوسط التقنية

³ استاذ مساعد: المعهد التقني – كوفة – جامعة الفرات الاوسط التقنية

⁴ استاذ دكتور: قسم تقنيات المختبرات الطبية – كلية الحكمة الجامعة – بغداد - العراق

⁵ استاذ مساعد: المعهد التقني – كوفة – جامعة الفرات الاوسط التقنية