

## **Comparing Range of Motion Between Females and Males Within Intervertebral Levels of Lumbar Segment**

**Raed A.M Al-Mamoori, Hayder Saad Oleiwi & Amer Adil Abdulkareem**

Al-Mussaib Technical College, Al-Furat Al-Awsat Technical University, 51009 Babylon, Iraq

*com.ma.raaed@atu.edu.iq*

**ABSTRACT:** The biomechanical characteristics of the healthy human spine for both of female and male are critical to be understood thoroughly so that the treatment of spinal pathology can be advanced. However, the rotational ranges of motion (ROM) are not the same for each sex. Hence, our goal is to; investigate the motion of the lumbar spine segment, as measured by range of motion ROM for the models of both sexes. A process works when by a CT scan of a lumbar spine is transformed into a simulation model. The study examines human lumbar segments; (L1-S1) responding to a pure-moment loading in: flexion extension (FE); lateral bending (LB) and also axial torsion (AT). The greater FE ROM is found at the lower lumbar levels alongside the significantly greater ROM for female segments compared with males. On that subject, there is greater flexibility in the female segments compared to males, and its influence on ROM has been discovered. These differences may be justified by higher incidence of LBP in females including the time when they are pregnant, and this causes the disc and facet degeneration deference rate in female more than male. The simulate models also could provide a critical input for: implant design; biomechanical study design; and interpretation, not to mention to the development of diagnostic criteria.

**KEYWORDS:** Range of motion; flexion extension; lateral bending; axial torsion

### **INTRODUCTION**

Low back pain (LBP) has become prevalent as a common reason for primary care visits after the common cold. It is reported that the majority of adults will have been suffering from this condition at some point in their lives [1, 2]. The Sacroiliac joint and lumbar spine constitute the largest axial joints in the body, as well as the spine articulation with the pelvis that allows loads being transferred to pelvis and lower extremities [3, 4]. Sexual dimorphism exists in pelvis it is different between the male and female sacrum. The female sacrum tends to be wider, not even, less curved, and that it tilts backward more. Additionally, males tend to have a relatively long and narrow pelvis, with a longer and more conical pelvic cavity than those of females [5].

We need to really comprehend the bio mechanical features for the healthy human spinal column for both of female and male because the treatment of spinal pathology has to be taken up a notch. As there is a problem of scarce vivo, and in-vitro studies, the state is a challenge toward quantify definite biomechanical parameters. The finite element analysis which could be utilized by understands of the biomechanics of the spine is very advantageous. The rotational ranges of motion (ROM) differ from one gender to another. The range of difference appears in the main loading modes, as flexion extension case (FE); also lateral bending (LB) in addition to axial torsion (AT), whereas they make common variables as well as they make the ultimate popular metrics that concerns with the biomechanics activity of the vertebral column.

Descriptive statistics ROM in healthy vertebrae and their relations with the demographical variables, can well contribute to: the implantation design with biomechanical investigating design; interpretation; Moreover, diagnostic criteria development; as well as design of bio mechanical spine models. Most of in-vitro studies for biomechanical of the vertebral column weigh in osseoligamentous creations unchanged.

Despite the varying methods to measure the Sacroiliac and lumbar segment motion [6-9] it is important to look closely into the biomechanical differences between male and female with regard to ROM. Cadaver studies demand a very technical procedure, as the motions at segments are very low. Furthermore, quantifying stresses across the joint is not feasible. With that, experimentally validated finite element analysis approach would be the best approach where the ROM, stresses, and strains across the joint can be evaluated.

We are going in the direction of consider the motion at the lumbar spinal column segment for both of male and female models as per measured by range of motion ability ROM, then study the dependences proceeding sex difference at

intervertebral spine level at the lumbar segment using female and male specific finite element models.. It is hoped that in terms of their mobility and the possible pain sites, relevant parties can better understand the biomechanical differences in lumbar spine between female and male.

**MATERIALS AND METHODS**

CT scans developed a validated finite element models in lumbar spine for both of female and male. The models helped to simulate spine physiological motions. A comparison between male and female models was made, with regard to the range of motion ROM across the left and right lumbar segment.

The study focuses on human spine segments (from L1 to S1). As well as, response to: a pure-moment; loading in flexion and extension case (FE); (LB) lateral bending; in addition to axial torsion (AT). The analyses were done so that the differences between the female and male in each mode of loading can be detected at intervertebral levels. Table 2 includes the Material properties for the models of both genders.










**Female- Male Finite Element LumbarSpine Model**

Computer tomography (CT) images of a 57 years old female’s and male’s spine without any abnormalities, degeneration helped to reconstruct the female and male lumbar spine model. MIMICS software (Materialise, Leuven, Belgium) was utilized to create a 3D geometry of the bones and then intervertebral discs were made by having the space between each two vertebrae of the CT images filled in. Next, smoothing and meshing were carried out with the help of the Geomagic Studio software (Raindrop Geomagic Inc., USA) and the Hypermesh software (Altair Engineering, Inc., USA). Table 1 illustrates the male and female spine FE models. The finite element lumbar spine model [10, 11] previously developed and validated was used with the current models. Lumbar spine bones were modeled as trabecular cores surrounded by a cortical layer. The linear hexahedral element type was used for cortical and cancellous bones of vertebrae and intervertebral discs. The truss elements were used for ligamentous tissues. 118,417 elements were generated for the male model, while female model as a whole contained 437,792 elements.

**Material Properties**

The material properties used in the FE models were extracted from previous studies [10, 13] and summarized in table 2.

**Table 1.** FE models parts of spine, (a) FE model of female lumbar spine -femur (b) FE model of male lumbar spine – femur

		
	(a)	(b)
		
(NP) and (AF)	End plate	Intervertebral Disc
		
Cancellous Bone	Cortical bone	Vertebrae

**Table 2.** Material properties of male and female models

Component	Material Properties	Constitutive relation	Element Type	Reference
Vertebral cortical bone	E= 12000 GPa v = 0.3	Isotropic, elastic	8 Nodes brick element (C3D8)	Lindsey et al. [10]
Vertebral cancellous bone	E = 100 GPa v = 0.2	Isotropic, elastic	8 Nodes brick element (C3D8)	Lindsey et al. [10]
Annulus fibrosus (Male model)	C10 = 0.3448 D1 = 0.3	Hyperelastic, neo-Hookean	Rebar	Lindsey et al. [10]
Annulus fibrosus (Female model)	C10 = 0.035 K1 = 0.296 K2 = 65	Hyperelastic anisotropic (HGO)	8 Nodes brick element (C3D8)	Shahraki et al. [13]
Nucleus Pulposus	E = 1 GPa v = 0.499	Isotropic, elastic	8 Nodes brick element (C3D8)	Lindsey et al. [10]
Ligaments	Nonlinear stress-strain curves	Hypoelastic	Tension-only, truss elements (T3D2)	Lindsey et al. [10]
* E: Is Young's elasticity modulus; v: Poisson's ratio; C1 and C2: Material constant characterizing the deviatoric deformation of material.				

### The Mesh Convergence Analysis

The mesh convergence analysis was carried out on the segregated L1-L5 motion segment of the female model. An initial seed size was assigned and the model was subjected to 7.5N.m bending moment to simulate motions in all planes, prior to the measurement of ROM. The final element size was used to mesh the other segments of the model. ABAQUS 6.14 software (Materialise, Leuven, Belgium) was used for the simulation.


### Loading and Boundary Conditions

In all models, a 400 N compressive follower load was applied through wire elements right after the curvature of the lumbar segment. This is to simulate the effect of muscle forces and weight of the upper trunk. To simulate the physiological flexion, extension, lateral bending, and axial rotation, a 10 N.m bending moment was then applied at the superior surface of the L1 vertebrae. For constraining the models, femurs were set in all degrees of freedom as to avoid relative displacement [10, 11].

## RESULTS

### Model Validations


The data predicted for all physiological loadings fell within one standard deviation of the experimental data, with the exception of the right lateral bending and right axial rotation for the male data, Figure 1. The intact male model L1-S ROM was validated before under the same loading and with the same posture conditions. For consistency, the validation under loading condition was done for both models. For intact validation, the loading conditions of the cadaver study analysed by Lindsey et al. [14] had to undergo a simulation. This experiment was carried out for intact L4 to pelvis for both gender specimens under loading condition. A 7.5 Nm pure moment load was applied to the top endplate of L4 in order for various spinal motions to be simulated. The motion at the segment was then calculated for both right and left joints.




**Figure 1.** Validation results for the intact female and male lumbar segment-sacrum (right and left sides) at 7.5 N.m moment under loading condition. Experimental data was taken from Lindsey et al. [14].

Comparing between the female and male range of motion ROM


At vertebral levels, comparing between males and females, ROM was greater for females in a greater way rather than males at: FE; LB; also AT. ROM at LB is wont to be greater toward a central part of segment per L2 and 3, L3 and 4; moreover L4-5 hence ROM being considerably larger than both of L1-2; L5-S1. These comparisons can be referred to in Figure 2, Figure 3 and Figure 4.



**Figure 2.** Comparisons Range of Motion at Flexion- Extension (FE) within Intervertebral Level between females and males



**Figure 3.** Comparisons Range of Motion at Lateral Bending LB within Intervertebral Level between females and males



**Figure 4.** Comparisons Range of Motion at Axial torsion AT within Intervertebral Level between females and males


ROM was considerably higher for females rather than for their counterparts of FE. When comparing, females demonstrated significantly higher ROM in each level of FE. These comparisons can be seen in Figure 2. In FE, ROM of Female lumbar spine model ranged had a greater ROM than Male lumbar spine model at L1-L2 and L3-L4, respectively. In flexion, the ROM of Female was again larger at L (3-4), also in L (4-5). The biggest comparative differences were noticed at state of extension, in which segments of a Female had highlighted a larger rate of ROM at L5-S1 which is greater at L4-L5. The ROM is wont to growth downward the vertebral column in FE; L3 with L4 FE ROM are recorded to be significantly more than L1 and 2, so for L4 and 5, with L5 to S1 FE ROM were considerably greater from every other level. Flexion ROM lean towards to be greater in the direction of the center of the segment with L2 and L3, L3- 4 and L4-L5 ROM being significantly greater than both L1-L2 and L5-S1. A similarly in trending was establish with extension. Nevertheless, just L1-L2 was a lesser amount in a striking way than all other levels. In LB, ROM of Female specimens were greater than that of males at L3-4 and L4-5, respectively (as shown in figure 3). The higher comparative variances is evident at (AT), in which is the Female segments had shown a larger mean value of ROM at L5-S1 and at L4-5 (as shown in fig 4). The LB of ROM lean towards to be greater in the direction of the center of segment within L2 and 3, also L3 and 4 until L4 and 5. A similarly trend founded at AT case. However, only for L1 and 2 were noted to be lesser from whole other levels.

It is a fact that many studies have been conducted to define the effects of disc degeneration, then facet osteoarthritis happened on lumbar spine flexibility, and subsequently there are different forms of the effect of disc furthermore facet degeneration taking place on ROM among females and males. A number of factors may be the reasons for the variability for ROM values. It can be stated that the height, weight and age may contribute to the differences in ROM values.

The current study has analysed the difference between lumbar segment ROM of female and male in different motions. The presented data helped to address the many critical questions about the male and female lumbar segment's anatomical variation. The indication is that men would have a greater lever arm than women, as the former has stronger sacroiliac joints [15, 18]. This characteristic may explain why males are known to be less mobile.

## DISCUSSION

In the current study we have seen that the female segment is appeared to be further flexible than males, and there is evidence of their influence on ROM. There were high increasing in FE of ROM in the lower levels at L4 -L5 also L5 until S1. Furthermore, the maximum LB ROM was at L (2-3). The greater FE ROM found at the lower lumbar levels additional to significantly higher ROM at females segment as compared to a males. Hence, a varying in pattern was influencing of intervertebral disc, and facet degeneration on ROM among males and females which have been observed. The results seem likely that geometrical differences between intervertebral levels in additions to between males and females offer a significant contribution to the differences in this study as pointed in Fig 5 [19-22].



**Figure 5.** Percentage of differences between (Female than Male) range of motion

The influence of FSU dimensions on the ROM response has somehow altered the percentage of difference for the results between both sexes. The difference can to be justified by the greater dimension size of male segments and lack of correspondence evident in properties of noted material. Last but not least, there are some significant correlations between height, weight, gender and ROM. Despite their weakness, they can still explain the differences of the current result.

## CONCLUSION

In conclusion, higher range of motion has been found in the female lumbar segment compared to the male model at both sides of the lumbar joint. Also, the act of stabilizing the lumbar segment was performed by one of the strongest ligaments in the body, the sacroiliac. These differences may lead to higher incidence of LBP in females, even though they are carrying a baby. We must consider the important variances in lumbar ROM between female and male spinal column segments, and between the intervertebral levels in study design; to avoid biases in outcomes. a female spinal column have been known to have higher flexibility or mobility; stresses; pelvis ligament strains and loads, as a compared with a male. Furthermore, it is causing higher rate of stress through the joint, principally at the sacrum in a similar loading conditions. Hence, a well explanation about: why there is larger incidence of sacroiliac ache with pelvic stress fracture in body of females. The implications within the conclusions or outcomes could strictly be critical in the design of spinal implants, particularly those concentrating on maintaining or else restoring for a healthy ability for motion. The major correlations concerning weight; ROM and height, however which is not clear sense, so may still be able to explain the differences of the current result, also through the intra gender differences.

---

## REFERENCES

- [1] N. Weksler, G.J. Velan, M. Semionov, B. Gurevitch, M. Klein, V. Rozentsveig, et al. "The Role of Sacroiliac Joint Dysfunction in the Genesis of Low Back Pain: The Obvious Is Not Always Right". *Arch Orthop Trauma Surg*, vol. 127, no. 10, pp. 885-8, 2007.
- [2] J.W. Frymoyer. "Back pain and Sciatica". *N Engl J Med*, vol. 318, pp. 291-300, 1988.
- [3] E. Dietrichs. "Anatomy of the pelvic joints-a review". *Scand J Rheumatol*, vol. 88, pp. 4-6, 1991.
- [4] S.P. Cohen. "Sacroiliac Joint Pain: A Comprehensive Review of Anatomy, Diagnosis, and Treatment". *Anesthesia & Analgesia*, vol. 101, no. 5, pp. 1440-53, 2005.
- [5] A.M. Khattak, M. Younas, G.A. Khan, M.U. Rehman S. Nadeem, M. Safeer. "P-Separation Axioms In Supra Soft Topological Space". *Matrix Science Mathematic*, vol. 2, no. 2, pp. 07-10, 2018.
- [6] S. Sathishkumar, M. Kannan. "Design and Fatigue Analysis of Multi Cylinder Engine and Its Structural Components". *Acta Mechanica Malaysia*, vol. 2, no. 2, pp. 10-14, 2018.
- [7] U.K. Hassan, S. Kasim, R. Hassan, H. Mahdin, A.A. Ramli, M.F. Md Fudzee, M.A. Salamat. "Most Stationery Inventory Management System". *Acta Electronica Malaysia*, vol. 2, no. 2, pp. 10-13, 2018.
- [8] E. O. Omini, O. M. Akpang. "Cavity Detection Under Re-Enforced Concrete Floor Using Ground Penetration Radar". *Engineering Heritage Journal*, vol. 2, no. 2, pp. 11-18, 2018.
- [9] A. Vleeming, M.D. Schuenke, A.T. Masi, J.E. Carreiro, L. Danneels, F.H. Willard. "The sacroiliac joint: An overview of its anatomy, function and potential clinical implications".
- [10] E. Vlaanderen, N.E. Conza, C.J. Snijders, et al. "Low back pain, the stiffness of the sacroiliac joint: a new method using ultrasound". *Ultrasound Med Biol*, vol. 31, pp. 39-44, 2005.
- [11] H.A. Jacob, R.O. Kissling. "The mobility of the sacroiliac joints in healthy volunteers between 20 and 50 years of age". *Clinical Biomechanics*, vol. 10, no. 7, pp. 352-61, 1995.
- [12] B. Stuesson, A. Uden, A. Vleeming. "A radiostereometric analysis of movements of the sacroiliac joints during the standing hip flexion test". *Spine*, vol. 25, no. 3, pp. 364-8, 2000.
- [13] B. Stuesson, G. Selvik, A. Uden. "Movements of the sacroiliac joints: a roentgen stereophotogrammetric analysis". *Spine*, vol. 14, pp. 162-5, 1989.
- [14] D. Lindsey, A. Kiapour, S. Yerby, V. Goel. "Sacroiliac joint fusion minimally affects adjacent lumbar segment motion: a finite element study". *Int J Spine Surg*, vol. 9, pp. 64, 2015.
- [15] A.A. Ivanov, A. Kiapour, N.A. Ebraheim, V.K. Goel. "Lumbar Fusion Leads to Increases in Angular Motion and Stress Across Sacroiliac Joint". *Spine*, vol. 34, no. 5, 2009.
- [16] C. Bruna-Rosso, P.J. Arnoux, R.J. Bianco, et al. "Finite Element Analysis of Sacroiliac Joint Fixation under Compression Loads". *Int J Spine Surg*, vol. 10, pp. 16, 2016.
- [17] N.M. Shahraki, A. Fatemi, V.K. Goel, A. Agarwal. "On the use of biaxial properties in modeling annulus as a Holzapfel-Gasser-Ogden material". *Frontiers in bioengineering and biotechnology*, vol. 3, 2015.
- [18] D. Lindsey, L. Perez-Orribo, N. Rodriguez-Martinez, P.M. Reyes, A. Cable, G. Hickam, A. "Newcomb Evaluation of a minimally invasive procedure for sacroiliac joint fusion - an in vitro biomechanical analysis of initial and cycled properties". *MDER Medical Devices: Evidence and Research*, vol. 131, 2014.
- [19] E.R. Tischauer, M. Miller, I.M. Nathan. "Lordosimetry: a new technique for the measurement of postural

---

response to materials handling”. *Am Ind Hyg Assoc J*, vol. 1, pp. 1-12, 1973.

- [20] N. Bellamy, W. Parl, P.J. Rooney. “What do we know about the sacroiliac joint?” *Semin Arthritis Rheum*, vol. 12, pp. 282-313, 1983.
- [21] N. Zheng, L.G. Watson, K. Yong-Hing. “Biomechanical modelling of the human sacroiliac joint”. *Med Biol Eng Comput*, vol. 35, pp. 77–82, 1997.
- [22] D.J. Cook, M.S. Yeager, B. C. Cheng. “Range of Motion of the Intact Lumbar Segment: A Multivariate Study of 42 Lumbar Spines”. *The International Journal of Spine Surgery*, 2018. DOI: <https://doi.org/10.14444/2005>.





Ref:  
Date:

العدد / ٤٥٧٨ / ٢٧ / ١٧ / ١٤  
التاريخ / ٢٠٢٠ / ١١ / ١٩

الى / كليات ومعاهد الجامعة كافة / السيد العميد المحترم  
م/ توثيق رصانة جهة نشر بحث

- نهديكم اطيب التحيات.....
- اشارة الى مصادفة السيد رئيس الجامعة المحترم على محضر لجنة الرصانة العلمية للنشر في المجلات العلمية لمستويات كلاريفيت (تومسن رويترز سابقاً) وسكوبس لاغراض ترصين البحوث العلمية المشكلة بموجب الامر الجامعي ذي العدد ١٠٨٨/٢٧/٧/٢ في ٢٠٢٠/٢/٢، نرافق ريبطاً جدول بالبحوث كلا حسب تصنيفه:
- ١- الجدول رقم (١) ابتداءً من التسلسل (١- ا.م.د. فاضل عباس جمعه) وانتهاءً بالتسلسل (٦٢- د. مرتضى محمد جواد) بحوث موثقة منشورة في مجلات سكوبس.
  - ٢- الجدول رقم (٢) ابتداءً من التسلسل (١- م.م. امير جواد كاظم) وانتهاءً بالتسلسل (٢٩- ليلي صالح عبد الحسن) بحوث مقبولة للنشر او منشورة وغير متوفره اولين في مجلات مصنفة ضمن تصنيف سكوبس.
  - ٣- الجدول رقم (٣) ابتداءً من التسلسل (١- د. ايمان جواد كاظم) وانتهاءً بالتسلسل (٤٤- زهراء حميد عودة القريشي) بحوث منشورة او مقبولة للنشر في مجلات مصنفة ضمن تصنيف سكوبس ويمكن توثيق رصانتها استناداً الى موافقة السيد رئيس الجامعة المحترم بتاريخ ٢٠٢٠/٤/١٠ على توثيق رصانة البحوث كون الموعد النهائي لتحديث قاعدة بيانات سكوبس لسنة ٢٠٢٠ يكون بتاريخ ٢٠٢٠/٥/٣٠، وان البحوث المرفقة في الجدول (٣) تعتبر منشورة او مقبولة للنشر في مستوعات سكوبس لسنة ٢٠١٩.
  - ٤- الجدول رقم (٤) ذات تسلسل (١- رباب عدنان فاضل و ٢- اسراء عبد الجليل حسين)، يتم توثيق رصانة البحوث بعد التأكد من قبول النشر قبل ٢٠١٩/٢/٣، اما بالنسبة للتسلسل (٣- د. اسعد سموم دغل) بحث قبول النشر بعد ٢٠١٩/٢/٣ ولم يتم توثيق رصانته حسب اعمام الوزارة.
  - ٥- الجدول رقم (٥) ابتداءً من التسلسل (١- ساهرة عايد عبد الصاحب) وانتهاءً بالتسلسل (٢- ساهرة عايد عبد الصاحب)، بحوث منشورة في مجلات غير مصنفة وليست تصنيف سكوبس او كانت ضمن تصنيف سكوبس وخرجت منه.

للتفضل بالعلم .... مع فائق الشكر والتقدير

المرفقات:

- الجدول رقم (١، ٢، ٣، ٤، ٥).

ا.م.د. احمد غانم وداي  
مساعد رئيس الجامعة للشؤون العلمية  
٢٠٢٠/٤/


نسخه منه الى:

- ❖ مكتب السيد رئيس الجامعة المحترم... لتفضل سيادته بالاطلاع... مع التقدير.
- ❖ شعبة الترقيات العلمية... لتفضل بالاطلاع... مع التقدير.
- ❖ قسم ضمان الجودة والاداء الجامعي / السيد مقرر لجنة توثيق البحوث العلمية... مع الاوليات.
- ❖ ملفه الكتب الصادرة.

Scopus 2009 to 2018	7 (4.3) (2018)	International Journal of Engineering & Technology	Numerical analysis of the informational parameters of the magnetic fields in the area of the defects of wheel pairs axes	د. جاسم محمد جاسم	الكلية التقنية / المسيب	9
Scopus 2013 to Present	Vol. 24, Issue 03, 2020	International Journal of Psychosocial Rehabilitation	Patterns of the Organizational Structure and its Impact in Meeting the Requirements of the Learn Organization: Applied Research in the Distribution Company of Oil Products in Basra	امير جواد كاظم	الكلية التقنية / المسيب	10
Scopus 2014 to Present	43 - 2 - 2020	Journal of Mechanical Engineering Research and Developments	Comparing Range of Motion Between Females and Males Within Intervertebral Levels of Lumbar Segment	راند علي محمد يوسف	الكلية التقنية / المسيب	11
Scopus 2010 to 2019	12(2) 2019	Journal of Biopesticides	Control of the causes of the damping off disease on pepper by some biological agents in Babylon province-Iraq	عهد عبد علي هادي مطلوب	الكلية التقنية / المسيب	12
Scopus Proceeding conference	745 (2020) 012003	IOP Conf. Series: Materials Science and Engineering	Optimal dual design based on capacitor placement and reconfiguration techniques for loss reduction and voltage enhancement	د.وليد خالد شاكر الجبوري	الكلية التقنية / المسيب	13
Scopus Proceeding conference	2019	5th Scientific International Conference Najaf, SICN 2019	Optimal Distribution System Reconfiguration Using Qualified Binary Particle Swarm Optimization Algorithm	د.وليد خالد شاكر الجبوري	الكلية التقنية / المسيب	14
Scopus Proceeding conference	2018	1st International Scientific Conference of Engineering Sciences - 3rd Scientific Conference of Engineering Science	Dynamic evaluation of girder cross-sectional shapes of bridges	ا.م.د. علي فاضل ناصر	الكلية التقنية / المسيب	15
Scopus Proceeding conference 2005 to present	Volume 1294, Issue 9	Journal of Physics: Conference Series,	Effect of locally manufactured implement on energy utilization efficiency and operating costs	ا.م.د. فراس جمعة	الكلية التقنية / المسيب	16
Scopus 2005 to Present	1372 (2019) 012022	Journal of Physics: Conference Series	Intelligent Fuzzy PD+I Controller with Stabilizer for Nano Satellite	محمد حساب مهدي	الكلية التقنية / الادارية / كوفة	17



02.2020.81.88



# COMPARING RANGE OF MOTION BETWEEN FEMALES AND MALES WITHIN INTERVERTEBRAL LEVELS OF LUMBAR SEGMENT

**Author(s):**

Raed A.M Al-Mamoori, Hayder Saad Oleiwi & Amer Adil Abdulkareem

**Affiliation(s):**

Al-Mussaib Technical College, Al-Furat Al-Awsat Technical University, 51009 Babylon, Iraq

*\*Corresponding Author Email: [com.ma.raaed@atu.edu.iq](mailto:com.ma.raaed@atu.edu.iq)*

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

[DOWNLOAD PDF](#)